

BCL2 Break Apart FISH Probe Kit

Introduction

The BCL2 Break Apart FISH Probe Kit is designed to detect rearrangements in the human BCL2 gene located on chromosome band 18q21.33. In addition to revealing breaks, which can lead to translocation of parts of the gene, inversion, or its fusion to other genes, the probe set can also be used to identify other BCL2 aberrations such as deletions or amplifications. Rearrangements and abnormal expression of the BCL2 gene – also known as Bcl-2 or PPP1R50 – is routinely found in follicular lymphoma but also occurs in many other hematological and solid cancer types. One particularly common rearrangement is a reciprocal translocation to the IGH@ locus on chromosome 14.

Intended Use

To detect rearrangements in the human *BCL2* gene located on chromosome band 18q21.33.


Cont.

Color

LSP BCL2 5' FISH Probe
LSP BCL2 3' FISH Probe

CytoOrange
CytoGreen

Probe Design



Not to Scale

LSP BCL2 5' FISH Probe covers sequences adjacent to the 5' (start) portion of the *BCL2* gene. LSP BCL2 3' FISH Probe covers some sequences downstream (3' end) of the gene. The two probes are flanking sequences across the *BCL2* gene in which variable breakpoints have been observed.

Cat. No.

Volume

CT-PAC206-10-OG

10 Tests (100 µL)

Signal Pattern Interpretation

Normal Patterns

2F*

Abnormal Patterns

Other Patterns

*Overlapping orange and green signals can appear as yellow.

- 1) Chao DT & Korsmeyer SJ. *Annu Rev Immunol.* 16:395-419 (1998).
- 2) Willis TG & Dyer MJ. *Blood.* 96(3):808-22 (2000).
- 3) Ngan BY, et al. *N Engl J Med.* 318(25):1638-44 (1988).
- 4) Einerson RR, et al. *Am J Clin Pathol.* 124(3):421-9 (2005).
- 5) Vaandrager JW, et al. *Genes Chromosomes Cancer.* 27(1):85-94 (2000).

* CE IVD only available in certain countries. All other countries are either ASR or RUO. Please contact your local dealer or our headquarters for more information.

DCN032

© CytoTest Inc.



CytoTest Inc.
1395 Piccard Drive, Suite 308
Rockville, MD 20850, USA