

## CRLF2 Break Apart FISH Probe Kit

### Introduction

The CRLF2 Break Apart FISH Probe Kit is designed to detect rearrangements in the human CRLF2 gene located on chromosome bands Xp22.33 and Yp11.2. In addition to revealing breaks, which can lead to translocation of parts of the gene, inversion, or its fusion to other genes, the probe set can also be used to identify other CRLF2 aberrations such as deletions or amplifications. Rearrangements and abnormal expression of the CRLF2 gene – also known as CRL2, TSLPR or CRLF2Y – have been observed in adult and pediatric acute lymphoblastic leukemia (ALL) and various other malignancies.

### Intended Use

To detect rearrangements in the human *CRLF2* gene located on chromosome bands Xp22.33 and Yp11.2.


### Cont.

### Color

LSP CRLF2 5' FISH Probe  
LSP CRLF2 3' FISH Probe

CytoOrange  
CytoGreen

### Probe Design



The human *CRLF2* gene locates on both chromosome X and Y. LSP CRLF2 5' FISH Probe covers some genomic sequences adjacent to the 5' end of the *CRLF2* gene. LSP CRLF2 3' FISH Probe covers the 3' part as well as sequences downstream of the gene. The two probes are flanking sequences across the *CRLF2* gene in which variable breakpoints have been observed.

### Cat. No.

### Volume

CT-PAC114-10-OG

10 Tests (100 µL)

### Signal Pattern Interpretation

#### Normal Patterns

2F\*

#### Abnormal Patterns

Other Patterns

\*Overlapping orange and green signals can appear as yellow.

1) O'Connor C. *Nature Education*. 1(1):171 (2008).  
2) Tsuchiya KD. *Clin Lab Med*. 31(4):525-42, vii-viii (2011).  
3) Ried T, et al. *Hum Mol Genet*. 7(10):1619-26 (1998).  
4) Tonoizuka Y, et al. *Cytogenet Cell Genet*. 93(1-2):23-5 (2001).  
5) Rand V, et al. *Blood*. 117(25):6848-55 (2011).

\* CE IVD only available in certain countries. All other countries are either ASR or RUO. Please contact your local dealer or our headquarters for more information.